

Chiral phase in fermionic measure and the resolution of the strong CP problem

P. Mitra*

Saha Institute of Nuclear Physics,
1/AF Bidhannagar,
Calcutta 700064, India

hep-th/0104186

Abstract

The fermionic measure in the functional integral of a gauge theory suffers from an ambiguity in the form of a chiral phase. By fixing it, one is led once again to the conclusion that a chiral phase in the quark mass term of QCD has no effect and cannot cause CP violation.

At present, a popular approach to anomalies is in the functional integral formalism. Here an integration is carried out over different field configurations. The integrand involves the classical action which possesses all classical symmetries, but the measure of integration may break some of these [1], and lead to anomalies, which can be calculated if a regularization is employed. The measure of integration over fermions has been extensively studied in connection with chiral anomalies, but as will be demonstrated here, there is still a certain ambiguity left. This ambiguity relates to the case when the fermion mass is non-vanishing, and is practically relevant in the important context of QCD. In the standard model, the parity-violating electroweak sector gives rise to an unknown chiral phase in the quark mass term of QCD, and this has been at the basis of the strong CP problem. The ambiguity in the fermion measure would imply an ambiguity in the amount of CP violation caused by the chiral phase in the fermion mass term. The natural way of fixing the ambiguity makes the CP violation disappear. This is in consonance with the results obtained by explicit regularization of the action, namely the generalized Pauli-Villars regularization [2] and lattice regularizations [3]. This reconfirms the basic idea behind the solution of the strong CP problem in the more formal framework in which anomalies arise

*mitra@tnp.saha.ernet.in

primarily from the measure and the regularization is relegated to the background. Indeed, this investigation was motivated by the fact that the earlier discussions resolving the strong CP problem were restricted to regularized actions and trivial measures, whereas many of the misconceptions that made up the strong CP problem arose in analyses using nontrivial measures instead of regularizations.

For a theory like quantum chromodynamics, with fermions interacting vectorially with gauge fields, the euclidean action may be written as

$$S = \int \bar{\psi}(\gamma^\mu D_\mu - m)\psi + S_g, \quad (1)$$

where D is the covariant derivative, hermitian γ -matrices are used and S_g stands for the gauge field action. The fermionic integration is supposed to be over $\psi, \bar{\psi}$, but in the context of anomalies, these are conventionally expanded in orthonormal eigenfunctions of $\gamma^\mu D_\mu$ [1],

$$\psi = \sum_n a_n \phi_n, \quad \bar{\psi} = \sum_n \bar{a}_n \phi_n^\dagger, \quad (2)$$

and the expansion coefficients $a_n, \bar{a}_n$, which are Grassmann variables, are integrated over. The functional integral is

$$Z = \int \mathcal{D}A \prod_n \int da_n \prod_n \int d\bar{a}_n e^{-S}. \quad (3)$$

A chiral transformation of the fermion fields

$$\psi \rightarrow e^{i\alpha\gamma^5} \psi, \quad \bar{\psi} \rightarrow \bar{\psi} e^{i\alpha\gamma^5}, \quad (4)$$

causes $a_n, \bar{a}_n$ to change,

$$\begin{aligned} a_n &\rightarrow \sum_m \int \phi_n^\dagger e^{i\alpha\gamma^5} \phi_m a_m, \\ \bar{a}_n &\rightarrow \sum_m \bar{a}_m \int \phi_m^\dagger e^{i\alpha\gamma^5} \phi_n, \end{aligned} \quad (5)$$

and there is a nontrivial Jacobian which can be calculated with an appropriate regularization. It is this Jacobian which is responsible for the axial anomaly in this approach [1] because the action is chirally invariant when $m = 0$. When $m \neq 0$, the divergence of the axial current gets a contribution from the explicit breaking due to this mass in addition to the anomaly.

We are interested here in the situation where the fermion mass term has a chiral phase. Let us write the action as

$$S_{\theta'} = \int \bar{\psi}(\gamma^\mu D_\mu - m e^{i\theta'\gamma^5})\psi + S_g, \quad (6)$$

where the prime in the symbol θ' is used to distinguish it from the coefficient of the $F\tilde{F}$ term in QCD. The corresponding functional integral will be given by an equation like (3), but what are the corresponding $a_n, \bar{a}_n$? One may naïvely follow [1] and use (2). In the presence of the chiral phase in the mass term, we shall distinguish these expansion coefficients by the superscript 0, writing

$$\psi = \sum_n a_n^0 \phi_n, \quad \bar{\psi} = \sum_n \bar{a}_n^0 \phi_n^\dagger, \quad (7)$$

and the functional integral is

$$Z_0 = \int \mathcal{D}A \prod_n \int da_n^0 \prod_n \int d\bar{a}_n^0 e^{-S_{\theta'}}. \quad (8)$$

One may, instead, try to *derive* a functional integral in the presence of a chiral phase in the fermion mass term from the simple case where there is no such phase. $S_{\theta'}$ is obtained from S by a chiral transformation (4) with $\alpha = \theta'/2$, so the expansion (2) gets altered to

$$\psi = e^{-i\theta'\gamma^5/2} \sum_n a_n^{\theta'} \phi_n, \quad \bar{\psi} = \sum_n \bar{a}_n^{\theta'} \phi_n^\dagger e^{-i\theta'\gamma^5/2}. \quad (9)$$

The chiral phase sticking out on the right hand side may look unfamiliar and strange, but it has to be remembered that the basis of expansion is not *a priori* fixed [4], and the phase can be thought of as causing a change of basis. Correspondingly

$$Z_{\theta'} = \int \mathcal{D}A \prod_n \int da_n^{\theta'} \prod_n \int d\bar{a}_n^{\theta'} e^{-S_{\theta'}}. \quad (10)$$

This integral is different from the previous one because of the anomaly: A redefinition of the expansion coefficients along the lines of (5), followed by a calculation of the Jacobian can be used to convert $a_n^{\theta'}, \bar{a}_n^{\theta'}$ to $a_n^0, \bar{a}_n^0$, but an $F\tilde{F}$ term gets added to the action, showing the difference between $Z_0, Z_{\theta'}$. Z_0 violates CP, while $Z_{\theta'}$ does not, because it is equivalent to the case with no chiral phase in either action or measure.

This already shows the ambiguity in the fermion measure, but one can be even more general. Note that Z_0 has been defined by postulating the standard measure in the presence of θ' , and $Z_{\theta'}$ has been derived by falling back on the standard measure only in the absence of any chiral phase in the mass term. There is a continuum of possibilities in between: one can start with a mass term having some chiral phase different from zero and θ' , and postulate the standard measure in that situation. Then one makes a chiral transformation to reach θ' in the mass term, but this chiral transformation alters the expansion of the fields. Thus, in general,

$$\psi = e^{-i\beta\gamma^5/2} \sum_n a_n^\beta \phi_n, \quad \bar{\psi} = \sum_n \bar{a}_n^\beta \phi_n^\dagger e^{-i\beta\gamma^5/2}, \quad (11)$$

and the functional integral is

$$Z_\beta = \int \mathcal{D}A \prod_n \int da_n^\beta \prod_n \int d\bar{a}_n^\beta e^{-S_{\theta'}}. \quad (12)$$

Here β is a real parameter. $Z_0, Z_{\theta'}$ correspond to the special cases $\beta = 0, \theta'$ respectively. This functional integral depends on β because of the nontrivial β -dependent Jacobian which would be involved in removing the chiral phase from the expansion by redefining the expansion coefficients: one would get a β -dependent $F\tilde{F}$ term added to the action. This β -dependence in the expansion, which can be converted to a β -dependence in the action, is the ambiguity referred to above.

This new parameter β may appear unnecessary because it has been overlooked earlier, but it can be nonzero, and if it is, it changes all old discussions of strong CP physics. The standard picture used to be that there are two parameters: one in the chiral phase of the fermion mass term, and the other in the $F\tilde{F}$ term hidden here in S_g . By chiral transformations one would then try to show that physical quantities depend on the algebraic combination of these two phases. What we are pointing out here is that the measure accommodates a third phase, and it is the algebraic combination of these *three* phases that is relevant for physics. A third phase also arises if explicit regularizations are used as in [2, 3] and has the same significance in the context of strong CP physics. It may be recalled that that phase gets fixed by the desire to maintain the classically conserved parity (see below) instead of running into regularization artefacts.

One may ask whether the anomaly depends on the parameter β . It is easy to see that even in the presence of a chiral phase in the expansion as in (11), a chiral transformation (4) leads to the change (5) of the expansion coefficients, and the anomaly is standard.

Can one then choose between the different values of β ? A choice is already suggested by the way the alternatives were introduced above. Z_0 and the more general Z_β for $\beta \neq \theta'$ were obtained by postulating the standard measure in an *unknown* domain, involving the presence of a chiral phase, while $Z_{\theta'}$, corresponding to $\beta = \theta'$ was derived from the standard measure in the *known* domain where there is no chiral phase, which can probably be trusted more. In view of the conflict, this may be the functional integral to be adopted.

Another way of choosing the parameter β is through the consideration of parity, as in [2, 3]. As explained there, the classical theory with a chiral phase in the mass term does *not* violate parity, in spite of appearances to the contrary. One simply has to redefine the parity operation for fermions. As it is, the parity operation for fermions in the absence of a chiral phase in the mass term involves a γ^0 . In the presence of the chiral phase θ' , this is altered and the appropriate parity operation is

$$\begin{aligned} \bar{\psi}(x_0, \vec{x}) &\rightarrow \bar{\psi}(x_0, -\vec{x}) e^{i\theta' \gamma^5} \gamma^0 \\ \psi(x_0, \vec{x}) &\rightarrow \gamma^0 e^{i\theta' \gamma^5} \psi(x_0, -\vec{x}). \end{aligned} \quad (13)$$

This change is justified by the fact that the changed transformation, together with the standard parity operation for gauge fields, leave $S_{\theta'}$ invariant, except to the extent that there is an $F\tilde{F}$ term in S_g , which explicitly violates parity. As $S_{\theta'}$ has this symmetry, it is natural to try to see if the fermion measure can be arranged to have the same symmetry. If not, it would be a case of an anomaly. Fortunately, there is no parity anomaly: the measure preserves the symmetry, *i.e.* the above transformation leaves $a_n^\beta, \bar{a}_n^\beta$ in (11) invariant if and only if $\beta = \theta'$. This can be seen by noting that under the parity operation,

$$\phi_n(x_0, \vec{x}) \rightarrow \gamma^0 \phi_n(x_0, -\vec{x}), \quad (14)$$

so that

$$\begin{aligned} \phi_n^\dagger(x_0, \vec{x}) e^{-i\beta\gamma^5/2} &\rightarrow \phi_n^\dagger(x_0, -\vec{x}) e^{-i\beta\gamma^5/2} e^{i\beta\gamma^5} \gamma^0, \\ e^{-i\beta\gamma^5/2} \phi_n(x_0, \vec{x}) &\rightarrow \gamma^0 e^{i\beta\gamma^5} e^{-i\beta\gamma^5/2} \phi_n(x_0, -\vec{x}). \end{aligned} \quad (15)$$

This confirms that one has to take $\beta = \theta'$. If one uses $\beta \neq \theta'$, and in particular the value zero, one is guilty of violating parity by a deliberate choice of measure when a parity conserving measure is available. Parity violation is then an artefact of the measure.

Unfortunately, the chiral phase θ' in the expansion has been overlooked in the literature and this has caused immense confusion. In particular, if a chiral transformation is carried out to remove θ' from the action, it automatically goes away from the expansion relations connecting the fermion fields with the eigenfunctions of the Dirac operator. No Jacobian needs to be calculated, and no $F\tilde{F}$ term gets generated in the action [5]. This means that the chiral phase has no effect, in agreement with the recent resolution of the strong CP problem [2, 3] through similar demonstrations that a chiral phase in the quark mass term does not violate CP. It is of interest to note that in all cases a chiral phase had to be identified to absorb the chiral phase in the mass term. In the other cases, this absorbing chiral phase appeared in the regulators, while in the present one it appears in the expansion or equivalently in the measure.

The problem studied here arose out of discussions with Haridas Banerjee on the strong CP problem.

References

- [1] K. Fujikawa, Phys. Rev. **D21** (1980) 2848
- [2] H. Banerjee, D. Chatterjee and P. Mitra, hep-ph/0012284
- [3] P. Mitra, hep-lat/0102008
- [4] *See, e.g.*, S. D. Joglekar and G. Saini, Ann. Phys. **229** (1994) 76
- [5] V. Baluni, Phys. Rev. **D19** (1979) 2227